

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Long-Term Air Pollution and Traffic Noise Exposures and Mild Cognitive Impairment in Older Adults: A Cross-Sectional Analysis of the Heinz Nixdorf Recall Study

Lilian Tzivian, Martha Dlugaj, Angela Winkler, Gudrun Weinmayr, Frauke Hennig, Kateryna B. Fuks, Mohammad Vossoughi, Tamara Schikowski, Christian Weimar, Raimund Erbel, Karl-Heinz Jöckel, Susanne Moebus, and Barbara Hoffmann, on behalf of the Heinz Nixdorf Recall Study Investigative Group

Table of Contents

Table S1. Spearman correlations between air pollution (ESCAPE-LUR) and noise variables (r_s)

Table S2. Association of noise with MCI, OR (95% CI)

Table S3. Association between L_{DEN} (categorical) and overall MCI

Figure S1. Effect modification of association between exposures and overall MCI. Panel A: Associations of $PM_{2.5}$ (per IQR) with overall MCI. Panel B: Associations of L_{DEN} (per 10 dB(A)) with overall MCI. Main model adjusted for age, sex, SES, alcohol consumption, smoking status, self-reported ETS, any regular physical activity and BMI.

Table S1. Spearman correlations* between air pollution (ESCAPE-LUR) and noise variables (r_s)

| Variable | PM_{2.5} ($\mu\text{g}/\text{m}^3$) | PM_{2.5} absorbance ($10^{-5}/\text{m}$) | PMcoarse ($\mu\text{g}/\text{m}^3$) | PM₁₀ ($\mu\text{g}/\text{m}^3$) | NO₂ ($\mu\text{g}/\text{m}^3$) | NOx ($\mu\text{g}/\text{m}^3$) | Traffic load at major roads (veh*m/d) | L_{DEN} (dB(A)) |
|--|---|--|---|--|---|--|--|--------------------------------|
| PM _{2.5} absorbance ($10^{-5}/\text{m}$) | 0.89 | | | | | | | |
| PMcoarse ($\mu\text{g}/\text{m}^3$) | 0.69 | 0.74 | | | | | | |
| PM ₁₀ ($\mu\text{g}/\text{m}^3$) | 0.88 | 0.89 | 0.70 | | | | | |
| NO ₂ ($\mu\text{g}/\text{m}^3$) | 0.65 | 0.62 | 0.46 | 0.54 | | | | |
| NOx ($\mu\text{g}/\text{m}^3$) | 0.63 | 0.52 | 0.42 | 0.51 | 0.93 | | | |
| Traffic load at major roads (veh*m/d) | 0.20 | 0.39 | 0.20 | 0.20 | 0.53 | 0.34 | | |
| L _{DEN} (dB(A)) | 0.30 | 0.48 | 0.32 | 0.31 | 0.37 | 0.29 | 0.59 | |
| L _{NIGHT} (dB(A)) | 0.30 | 0.48 | 0.32 | 0.32 | 0.36 | 0.28 | 0.57 | 0.99 |

*All the correlations are at 0.001 significance level

Table S2. Association of noise with MCI^a, OR (95% CI)

| | L_{DEN} continuous (per IQR=14.2 dB(A)) | L_{NIGHT} continuous (per IQR=13.6 dB(A)) | L_{DEN} (threshold 65 dB(A), per 10 dB(A)) | L_{NIGHT} (threshold 50 dB(A), per 10 dB(A)) |
|------------------|--|--|---|---|
| Overall MCI | 1.10 (0.94, 1.28) | 1.09 (0.94,1.27) | 1.89 (1.10, 3.25) | 1.35 (1.00, 1.82) |
| Amnestic MCI | 1.15 (0.94, 1.40) | 1.13 (0.93, 1.38) | 2.39 (1.28, 4.45) | 1.47 (1.02, 2.12) |
| Non-amnestic MCI | 1.06 (0.87, 1.30) | 1.06 (0.87, 1.30) | 1.34 (0.60, 2.98) | 1.22 (0.81, 1.85) |


^aAdjusted for age, sex, SES, alcohol consumption, smoking status, self-reported ETS, any regular physical activity, BMI

Table S3. Association between L_{DEN} (categorical) and overall MCI^a

| L_{DEN}, dB(A) | OR (95% CI) |
|-------------------------------|--------------------|
| < 45 | Reference |
| 45 - 55 | 0.98 (0.73,1.32) |
| 55 - 65 | 1.08 (0.78,1.49) |
| 65 - 75 | 1.21 (0.84,1.74) |
| > 75 | 1.77 (0.61,5.07) |

^aAdjusted for age, sex, SES, alcohol consumption, smoking status, self-reported ETS, any regular physical activity, BMI

A.



B.




Figure S1. Effect modification of association between exposures and overall MCI. Panel A: Associations of PM_{2.5} (per IQR) with overall MCI. Panel B: Associations of L_{DEN} (per 10 dB(A)) with overall MCI. Main model adjusted for age, sex, SES, alcohol consumption, smoking status, self-reported ETS, any regular physical activity and BMI.